	jc957
3,712	U.S.
_	-

Please type a plus sign in this box: +

PTO/SB/05 (12/97)

Approved for use through 9/30/00. OMB 0651-0032
Patent and Trademark Office, US DEPARTMENT OF COMMERCE

Patent and Trademark Office, US DEPARTMENT OF COMMERCE Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

UTILITY PATENT APPLICATION TRANSMITTAL

Attorney Docket No. | 014938.0003 | Total Pages | 1

First Named Inventor or Application Identifier

Chaoying Zhao

Express Mail Label No. | EL709376393US

(Only for new non-provisional applications under 37 C F.R. 1.53(b)

S	ee MF	APPLICATION ELEMENTS PEP chapter 600 concerning utility patent application	i contents	Al	DDRE	SS TO	Box Patent	ommissioner for Patents Application n, D.C. 20231	
1.		Fee Transmittal Form (Submit an original, and a duplicate for fee process) Specification [Total Pages 19] (preferred arrangement set forth below) - Descriptive title of the invention - Cross References to Related Applications - Statement Regarding Fed sponsored R & D - Reference to Microfiche Appendix]	. Nucleo	tide a	and/or A le, all ne Compu Paper (ecessary) iter Readable (Copy (identica	quence Submission	Jc912 U.S. PTO 09/713498
		- Background of the Invention			ACCO	OMPAN	NYING APPL	ICATION PARTS	
		- Brief Summary of the Invention	8	. 🗆	Assig	gnment	Papers (cover	sheet & documents)	
13		- Brief Description of the Drawings (if filed)	9	. 🗖	37 C	FR 3.73	(b) Statement	Power of Attorn	ey
₽J ₽ I		- Detailed Description	1	0.	Engli	ish Tran	slation Docum	nent (<i>if applicable</i>)	
i i		- Claim(s)	1	1.			Disclosure	Copies of IDS	
LU		- Abstract of the Disclosure			State	ment (II	DS)/PTO-1449	Citations	
And the state of t		Drawing(s) (35 USC 113) [Total Sheets	- 1			-	Amendment		
¥	\boxtimes	Oath or Declaration [Total Pag	ges 2] 1	3.			ipt Postcard (M	•	
LU E	a.	☐ Unexecuted (original or copy)			(Shoi	uld be s _i	pecifically iten	nized)	
the state white white the state of the state	b.	Copy from a prior application (37 CFR (for continuation/divisional with Box 17 comple [Note Box 5 Below] i. DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior app see 37 CFR 1.63(d)(2) and 1.33(b)	eted) 1 g 1	(if fo	State Copy oreign	n priorit	Status strity Document y is claimed)		
5.		Incorporation By Reference (useable if Box 4b		• • •					
		The entire disclosure of the prior application which a copy of the oath or declaration is su			. 				
		under Box 4b, is considered as being part of	~ ^		 				
		disclosure of the accompanying application a	and 1s	• • • •					
17	If a	hereby incorporated by reference therein. CONTINUING APPLICATION, check applications of the control of the con	propriate box o	ind supply	the r				
		Continuation Divisional	Continuat	ion-in-par	t (CII	P) of pri	or application	No:	
		18. C	ORRESPONI	DENCE A	DDR	RESS			
	Cus	tomer Number or Bar Code Label	20594			or		pondence address belo	w
NA.	ME DRE	ALVIN R. WIRTHLIN							
CIT			STATE	1			ZIP CODE		
CO	UNT	RY T	ELEPHONE				FAX		

Approved for use through 6/30/98. OMB 0651-0032

Patent and Trademark Office; US DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

						Complete if I	Known		
FEE TRANSMITTA	Application No.				N/A				
		Filing Date				November 15, 2000			
Note: Effective October 1, 19	97	First Named Inventor			itor	Chaoying Zhao			
.			o Art U			N/A			
Patent fees are subject to annual re	evision.		iner N			N/A			
TOTAL AMOUNT OF PAYMENT	\$355.00		ney Do				4938 0003		
		Numl	-			014938.0003			
		·							
METHOD OF PAYMENT (c)	heck one)				FEE C	ALCULATIO	ON (continued)		
 The Commissioner is hereby authorized indicated fees and credit any over payment. 			DDITIO Entity	ONAL FI Small	EES Entity				
Deposit 01-0657		Fee	Fee	Fee	Fee				
Account		Code	(\$)	Code	(\$)	Fe	ee Description	Fee	
Number	0 F 15	105	120	205		0 1 1.	C1 C	Paid	
Deposit Akın Gump Strauss Hau	er & Feld	105	130	205	65	_	filing fee or oath		
Account Name		127	50	227	25	cover sheet.	provisional filing fee or		
Charge Any Additional Charge the I	ssue Fee Set in	139	130	139	130	Non-English spe	ecification		
Fee Required Under 37 CFR 1.18	at the Mailing 37 CFR of Allowance	147	2,520	147	2,520		ication of SIR prior to		
Treat any future replay requiring an extensio incorporating a request therefor.	n of time as	112	920*	112	920*	Requesting publ Examiner action	lication of SIR prior to		
2. Payment Enclosed:		113	1,840*	113	1,840*	Requesting publ Examiner action	ication of SIR after to		
Check Money Order	Other	115	110	215	55		ply within first month		
FEE CALCULATION		116	390	216	195	Extension for re	ply within second month		
. FILING FEE		117	890	217	445	Extension for re	ply within third month		
Large Entity Small Entity		118	1,390	218	695	Extension for re	ply within fourth month		
Fee Fee Fee		128	1,890	228	945	Extension for re	ply within fifth month		
Code (\$) Fee Description	Fee Paid								
101 = 710 201 355 Utility filing fee		119	310	219	155	Notice of Appea			
106 = 320 206 160 Design filing fee	;	120	310	220	155	=	support of an appeal		
107 = 490 207 245 Plant filing fee		121	270	221	135	Request for oral	=		
108 1710 208 355 Reissue filing fe	i	138	1,510	138	1,510		ute a public use proceedir	ıg	
114 150 217 75 Provisional Filin	ig fee	140	110	240	55	Petition to reviv			
	~~~~	141	1,240	241	620	Petition to reviv			
SUBTOTAL (1)		142	1,240	242	620	Utility issue fee			
2. Claims Fee f Extra belo		143	440	243	220	Design issue fee	;		
Total Claims 7 -20= 0 X	9 = 0	144	600	244	300	Plant issue fee			
ndependent 1 -3 0 X	39 = 0	122	130	122	130	Petitions to the	Commissioner		
Claims  Multiple Dependent Claims  0 X	130 = 0	123	50	122	50	Patitions related	to provisional application	36	
Turupic Depondent Claims U A	150 - 0	123	240	123 126	240		nformation Disclosure Str	<u> </u>	
Large Entity Small Entity									
Fee Fee Fee Fee Code (\$) Code (\$) Fee Description		581	40	581	40	property (times	patent assignment per number of properties)		
103 18 203 9 Claims in excess 6	of 20	146	710	246	355	Filing a submiss (37 CFR 1.129(a	sion after final rejection		
102 80 202 40 Independent claim	ns in excess of 3	149	710	249	355	,	nal invention to be exami	ned	
104 270 204 135 Multiple depender	nt claim	Other fo	ee (specif	fy)		(2. 2211.12)(1	-,		
110 18 210 9 Reissue claims inc			ee (specif						
over original pate					P. D. J		CYIDTOTAL	(2)	
SUBTOTAL (2)	0	"Keduc	ed by Ba	sic Filing	Fee Paid		SUBTOTAL	, (3)	
SUBMITTED BY							Complete (1f	applicable)	
Typed or Printed Name Alvin R. Wirthlin							Reg. Number	40,267	
Signature 30 - Pl. li	+10.		I	Date			Deposit Account		

# of the state of th

### IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In repatent application of:

Chaoying Zhao

Serial No. N/A Group Art Unit: N/A

Filed: November 15, 2000 Examiner: N/A

For: NOVEL PHARMACEUTICAL COMPOSITIONS FOR TREATING AND

SAVING AND THE METHOD FOR THE PREPARATION THEREOF

Assistant Commissioner for Patents Washington, D.C. 20231

Sir:

### CERTIFICATE OF MAIL BY "EXPRESS MAIL"

"Express Mail" Mailing Label No. EL709376393US

Date of Deposit: November 15, 2000

I hereby certify that the attached patent application comprising the following documents is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. §1.10 in an envelope addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231 on the date of deposit indicated above.

- Utility Patent Application Transmittal;
- Fee Transmittal;
- Filing fee in the amount of \$355.00;
- 19 sheets of specification, claims and Abstract of the Disclosure;
- Declaration (unexecuted);
- Preliminary Amendment; and
- return postcard.

(Typed Name of Person Depositing Envelope in Express Mail Facility)

Alm Lovelace

Cianature

Akin, Gump, Strauss, Hauer & Feld, L.L.P.

P.O. Box 688

Dallas, TX 75313-0688 Phone: (214) 969-2800

#412456



Inventor One Given Name:: Chaoying

Family Name:: Zhao

Postal Address Line One:: Changhai Road 170-7-602

City:: Shanghai

Country:: P.R. China

Postal or Zip Code:: 200433 City of Residence:: Shanghai Country of Residence:: P.R. China Citizenship Country:: P.R. China

### CORRESPONDENCE INFORMATION

Correspondence Customer Number:: 20594

Fax One:: 214/969-4343

Electronic Mail One:: awirthlin@akingump.com

### APPLICATION INFORMATION

Title Line One:: NOVEL PHARMACEUTICAL COMPOSITIONS FOR TR Title Line Two:: EATING AND SAVING AND THE METHOD FOR THE

Title Line Three:: PREPARATION THEREOF

Formal Drawings?:: No Application Type:: Utility Docket Number:: 014938.0003

Secrecy Order in Parent Appl.?:: No

### REPRESENTATIVE INFORMATION

Representative Customer Number:: 20594

### PRIOR FOREIGN APPLICATIONS

Foreign Application One:: PCT/CN99/00055

Filing Date:: 04-16-1999

Country:: WIPO

Priority Claimed:: Yes

Foreign Application Two:: 98108902.X

Filing Date:: 05-15-1998 Country:: P.R. China Priority Claimed:: Yes

Source:: PrintEFS Version 1.0.1

### IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In repatent application of:

Chaoying Zhao

Serial No. N/A

Group Art Unit: N/A

Filed: November 15, 2000

Examiner: N/A

For:

NOVEL PHARMACEUTICAL COMPOSITIONS FOR TREATING AND SAVING AND THE METHOD FOR THE PREPARATION THEREOF

**Assistant Commissioner for Patents** 

Washington, DC 20231

### PRELIMINARY AMENDMENT

Sir:

Please amend the above-identified application as follows.

### In the Claims:

Please amend the claims as follows:

Claim 3, line 1, please delete "or 2";

Claim 4, line 1, please delete "or 2".

Respectfully submitted,

11/15/00

Alvin R. Wirthlin, Reg. No. 40, 267

Akin, Gump, Strauss, Hauer & Feld, L.L.P.

P.O. Box 688

Dallas, TX 75313-0688

(214) 969-2800

10

15

20

25

# Novel Pharmaceutical Compositions for Treating and Saving and the Method for the Preparation thereof

The present invention relates to a novel pharmaceutical composition for treating and saving the wounded, and to a method for the preparation thereof.

Nowadays blood transfusion and fluid infusion are the important measures to treat and save the wounded, especially the traumatic shock. Usually, the principle of transfusion is "to infuse component the patient is deficient in, to supply how much the patient needs". For example, when the patient mainly lost his blood, he should be transfused with blood, even though sometimes the patient needs to be transfused with blood from normal individuals. When the patient mainly lost plasma, plasma or plasma volume expander should be transfused to him; when the patient mainly lost intercellular fluid, physiological saline should be infused. In fact, the treating and saving measures of formulating physiological solutions on the basis of normal composition of body, or transfusing with blood from normal individuals to the patient with evident physiopathological changes is to treat the organism as mechanical device, therefore these measures often have following disadvantages:

(1) Blood transfusion: In general, the volume of blood transfusion closes to or exceeds the volume of blood lost. If a big amount of blood is required, the blood source will be difficult, the cost is expensive. The preparation and storage need certain conditions. In addition, before transfusion, some time should be taken for blood typing and cross match tests, and only the substitutes could be used for the individuals with rare blood types. Blood transfusion could result in production of anti--platelet antibodies and anti--leucocyte antibodies, as well as various hematogenic infectious diseases, for example, AIDS, hepatitis B, hepatitis C. etc.

10

15

20

25

- (2) Albumin infusion: There is a great demand, a great expense difficult source, complicated preparation method and certain requirements for, the method storage. After albumin infusion, it could effuse through capillaries, and couldn't be reabsorbed into vessel. Therefore interstitial edema will occur, and might result in pulmonary edema, renal failure, and cardiac insufficiency, by contrast enhance the mortality. Albumin infusion could result in evident decrease of  $\alpha_1$ ,  $\alpha_2$ ,  $\beta$ ,  $\gamma$ -globulins and fibrinogen, cause reduction of immunity, and affect function of blood coagulation.
- (3) Infusion of fluorocarbon as plasma substitute: It requires a large amount of the substitute and simultaneous inhalation of oxygen with high component pressure. Otherwise the demand of the organism is not met. The fluorocarbon as plasma substitute should be preserved in low temperature, and its transport is not convenient. For example, thirty minutes before infusion the patient should be injected with 10 mg of dexamethasone. The frozen injection for infusion should be thawed. Thus, the procedures are complicated. After infusion, the observation should be continued for 5 -- 10min. The infusion of fluorocarbon as plasma substitute could induce adverse reactions, such as anaphylaxis, hypotension, thrombocytopenia, hepatosplenomegaly, reduction of immunity, and abnormality of fibrinolysis system.
- (4) Infusion of balanced buffers, Its amount infused should be three times more than the volume of blood lost to maintain the blood pressure. 60-80% the solution infused could effuse out of the blood vessel, which results in tissue edema (for example, cerebral edema, pulmonary edema) and renal insufficiency, it could cause the difficulty for sequential therapy.
- (5) Infusion of physiological saline. Its amount infused should also be three times more than the volume of blood lost. Its efficacy is poorer than balanced buffers, and its adverse effects are more significant.

10

15

20

25

To solve the problems in blood transfusion and infusion, the skilled in the art had studied anti--shock therapy with hypertonic sodium chloride solution. For example, 7. 5% (w/v) NaC1 solution suggested by Velasco. But the hypertonic NaC1 solution has some toxicity to the organism.

Most investigators have proposed intravenous injection of hypertonic NaC1 solution for anti--shock therapy, but it usually leads to obvious complications, such as hypotension, rupture of blood cells induced by extra hypertonic solution, cardiac insufficiency, decreased renal function and disorders of nervous system.

Thus, it can be showed that there is a demand for novel anti--shock drugs to reverse the physiopathological condition of shock, in order to obtain time for sequential therapy after the emergency treatment, and to create opportunity with improved effect of treating and saving the wounded and patients, and with increased survival rate.

An object of the present invention is to provide a pharmaceutical composition with convenient source, less dosage, rapid and better efficacy, less side effects, wider uses, and without restriction by blood type as well as without special storage condition.

Another object of the present invention is to provide the method for the preparation of the said pharmaceutical composition.

The present invention proposes a new concept of liquid therapy for shock, based on three aspects of thinking. The first aspect in connection with the present unreasonable dosage regimen, following principle is adopted: "to infuse what component the patient needs, then to infuse how much the patient needs". Based on the physiopathological status of the patient with shock, there is a prior demand for the compound solution containing hypertonic sodium ion (or combination of various crystals, or combination of various crystals and various colloids, etc.) to preliminarily improve micro---

10

15

20

25

circulation, tissue perfusion, and hemodynamics immediately. Then in the light of practical demand the isosmotic solution or proper hypoosmotic solution or whole blood or concentrated red blood cell suspension is administered, in order to permit the latter infused solution better action when the patient's condition has improved preliminarily by the earlier hypertonic solution, and to remit over--dehydration of some cells caused possibly by the hypertonic solution infused earlier. The second aspect, in connection with the present unreasonable ratio of colloids and crystals in the transfusion for patient with shock prepared on basis of their normal physiological concentrations, in view of the fact that the property and volume--expanding ability of artificial colloids are different from albumin in blood, it is considered that the transfusion with suitable ratio of colloids and crystals, instead of human normal physiological proportion, should be administered on basis of the physiopathological status of patients. Thus medicine administration according to indications could reduce the dosage, increase the efficacy, and decrease the complications. The third aspect, in connection with a lot of inadequacy in the present anti--shock experimental studies (such as animal model, reasonable concentration and infusion rate of hyperosmotic solutions), a great number of experimental studies have been conducted, and met with success.

Based on the above three aspects of thinking form the theoretical researches and the clinical experiences. the particular embodiments of the present invention have been completed through the animal experiments and the clinical practice.

The present invention is achieved through the following embodiments. A pharmaceutical composition comprising 1.5—6.9% (w/v) of one or more substances selected from sodium chloride, sodium bicarbonate, potassium chloride, magnesium sulfate, calcium chloride, calcium gluconate, calcium

10

15

20

lactate, sodium lactate, sodium acetate and Tris (Hydroxy methyl) aminomethane; and 3--18%(w/v) of one or more substances selected form hydroxyethylstarch, dextran, carboxy methylstarch, polyvinyl- pyrrolidone (PVP), gelatin derivatives, condensed glucose, glucose, fructose, lactose, glycerin, xylitol, sodium alginate, N-2- hydroxypropylacrylamide, ethylene epoxide--polypropylene glycol, pectin, mannitol, and penta hydroxyethylstarch (Pentastarch) as well as the remainder of conventional injections, as long as sodium chloride is not less than 1. 5%(w/v), and the concentration of sodium ion is not more than that of in 6. 9% (w/v) sodium chloride solution or equivalent.

The preferred composition of the present invention consisting of 4.2±0.2g sodium chloride and 7.6±0.6g hydroxyethylstarch per 100ml.

In the composition, the said hydroxyethylstarch contains at least 10% hydroxyethylstarch with molecular weight of 25, 000--45, 000.

The said dextran has molecular weight of 40, 000--230, 000, carboxymethylstarch has molecular weight of 30, 000--80, 000, PVP has molecular weight of 5, 000--700, 000, condensed glucose has molecular weight of 8, 000--12, 000; sodium alginate has molecular weight of 20, 000--26, 000: pectin has molecular weight of 20, 000--40, 000: pentahydroxyethylstarch is the product of DuPont Company (Pentastarch) with molecular weight of 264, 000.

The said gelatin derivatives have molecular weight of 20, 000- 35, 000, and are selected from urea--conjugated gelatin, modified liquid gelatin, oxidized polygelatin and degraded gelatin poly--peptide.

Conventional injections are selected from water for injection, physiological saline, balanced buffers, glucose solution, sodium lactate solution, sodium acetate solution, Tris solution, and glucose and sodium chloride solution.

10

15

20

25

The composition of the present invention is prepared as following procedure: dissolving 3--18g of total amount of one or more substances selected from hydroxyethylstarch, dextran, carboxymethylstarch, PVP, gelatin derivatives, condensed glucose, glucose, fructose, lactose, glycerin, xylitol, N--2--hydroxypropylacrylamide, sodium alginate. ethylene epoxide-polypropylene glycol, pectin, mannitol, and pentahydroxyethylstarch in 100 ml of total volume of one injection or mixture of several injections selected from water for injection. physiological saline, balanced buffers, glucose solution, sodium lactate solution, sodium acetate solution, Tris solution, and glucose and sodium chloride solution; then adding 1. 5g sodium chloride and 0-5.4g of one or more substances selected from sodium chloride, sodium bicarbonate, potassium chloride, magnesium sulfate, calcium chloride, calcium gluconate, calcium lactate, sodium lactate, sodium acetate, and Tris; with the proportion described above, then mixing, and dissolving, to obtain the composition of the present invention.

The preferred technical embodiments are as follows: (i) preparation of hydroxyethylstarch: According to the proportion of 1:0. 8- 0. 875: 0. 04--0. 042 (w/v/v), corn starch or sorghum starch, and 95% ethanol, and 35--38% hydrochloric acid are mixed, the temperature is raised to 65--80 for hydrolysis of starch, 16% sodium hydroxide solution is added in proportion of 0. 6--0. 7: 1 (v/w) of stated solution versus starch. Then ethylene epoxide is added in proportion of 0. 35--0. 5: 1 (w/w) of ethylene epoxide versus starch, then the mixture is heated to 65--75 to cause starch hydroxyethylation. (ii) Formulation of the composition: Appropriate volume of water is added, to prepare 7.6±0.6%(w/v) hydroxyethylstarch solution. Suitable amount of activated charcoal is added to discolor the solution through adsorption. After filtration, the pH is adjusted to 5. 5--7, and the total of 4±0.2g of sodium chloride is added, suitable amount of activated charcoal is added again for

adsorption and discolor action. After filtrated through 0.8 m micro-porous filter, the preferred composition of the present invention is obtained.

The present invention is further illustrated in detail by the following examples.

Preparative example.

5

10

20

25

Preparation of hydroxyethylstarch.

100g Corn or sorghum starch are mixed with 87ml of 95% ethanol and 4. 2 ml of 35% hydrochloric acid. The temperature is raised to 70 to hydrolyze starch then 60ml at 16% sodium hydroxide solution is added, then 45g epoxyethane is added, and the mixture is heated to 70 to complete hydroxyethylation of starch. According to the formula and preparative method described above hydroxyethylstarch with molecular weight of 25, 000--45, 000 is obtained.

Example 1.

Prepare according to the following proportion:

hydroxyethylstarch

7.6g

sodium chloride

4.2g

water for in injection added to 100ml

7. 6g hydroxyethylstarch are dissolved in 100 ml of water for injection.
0. 5g of activated charcoal is added, and the mixture is heated at 90 for 15 min under stirring. After filtration through asbestos plate filter, 4. 2g sodium chloride (purity pharmaceuticals use) are added, and dissolved with stirring.
0.5g activated charcoal is added, and the mixture is heated at 90 for 10 min under stirring. After filtration through asbestos plate filter and 0. 8 m m i croporous filter, resulted filtrate is transferred into 250ml or 500-ml glass or plastic bottles (bags), after sealing the bottles or bags are 1.05 kg/cm² and 121-123 for 15--30min for sterilization, to obtain the pharmaceutical

10

15

20

25

composition of the present invention.

## Example 2.

Prepare according to the following proportion:

dextran	9g
hydroxyethylstarch	3g
sodium chloride	1.5g
sodium bicarbonate	3.4g
physiological saline	added to 100ml

Above--mentioned dextran (produced by Shanghai Glucose Factory), hydroxyethylstarch (prepared according to preparative example) are dissolved in physiological saline, and adsorbed and discolored with activated according to the method stated in Example 1. Then above--mentioned sodium chloride, sodium bicarbonate, are added in turn, and dissolved with stirring. There after the obtained solution is discolored, filtered, sterilized and filled, to obtain the pharmaceutical composition of the present invention.

# Example 3.

Prepare according to the following proportion:

polyvinyl--pyrrolidone (PVP) (produced by Bayer) 12g sodium chloride 2g sodium acetate 4g 10% glucose solution added to 100ml

According to the method described in Example 2 except that dextran and hydroxyethylstarch is replaced by PVP, sodium bicarbonate is replaced with sodium acetate, and physiological saline is replaced with glucose solution, the composition of the present invention is obtained.

# Example 4.

Prepare according to the following proportion:

sodium alginate (produced by Nanning

18g

Pharmaceutical Factory, Guangxi)

sodium chloride

1.5g

5 water for injection.

added to 100ml

According to the method described in Example 1 to prepare the above-mentioned formulation, thus obtaining the pharmaceutical composition of the present invention.

Example 5.

Prepare according to the following proportion:

pectin (produced by PLA No. 185 Hospital) 3g

Pentahydroxyethylstarch (produced by 4g

DuPont Company)

sodium chloride

4g

mannitol

7g

2% sodium lactate lution

added to 100ml

According to the method in described in Example 1, pectin Pentahydroethylstarch and mannitol are dissolved in sodium lactate solution, 20 then sodium chloride is added and dissolved.

Example 6.

Prepare according to the following proportion:

condensed glucose (produced by southwest 7g

No. 5 Pharmaceutical Factory Chongqing)

N-2-hydroxy propyl acrylamide 2g

sodium chloride

4.4g

water for injection

added to 100ml

Using the method in Example 1, the pharmaceutical composition of the present invention is prepared according to the above mentioned formula.

# Example 7.

5 Prepare according to the following proportion:

fructose (produced by Shanghai

5g

No. 2 Reagent Factory)

xylitol (produced by Liaoyang organic

4g

Chemical Plant)

sodium chloride

4.8g

water for injection

added to 100ml

Using the method in Example 1, the pharmaceutical composition of the present invention is prepared according to the above--mentioned formula.

Example 8.

Prepare according to the following proportion:

glycerin

2g

lactose (produced by Shanghai

5g

No. 2 chemistry Reagent Factory)

sodium chloride

6g

water for injection

added to 100ml

Using the method in Example 1, the pharmaceutical composition of the present invention is prepared according to the above-mentioned formula.

# 25 Test 1: Animal Experiment

From adult healthy hybrid dogs, regardless of sex, under local anesthesia, isolate femoral artery and femoral vein, then insert catheters respectively.

10

The arterial duct is connected to CF--II model monitor of cardiovascular function [Shanghai approval document number: Hu--Yao- Qi--Jian (Zhun)--97—221103] to monitor cardiovascular status. Bleed these dogs to monitor cardiovascular status. Bleed these dogs to average arterial pressure (MAP) of 40--50 mmHg for a period of about 15 min. Maintain this blood pressure level for 1 hour, then infuse the product prepared in Example 1 at the dose of 8 ml/kg.

Monitor the cardiovascular function and urine volume over 4 hours after infusion. In the following tables, the blood pressure and other indexes are expressed as percentage of their basal levels respectively, the unit of urine volume is ml/kg body weight/h.

Comparison between the composition of the invention and whole blood in equal volume of recovery of Cardiovascular function in dogs with shock. [unit: % compared with respective basal level] Table 1

			afi	after transfusion		
	i	30 min	1h	2h	3h	4h
Systolic pressure	composition of the invention	75±4	76±4	77±4	78±5	79±5
	equal volume of whole blood	67±5	9∓69	73±4	73±4	77±7
•		(p<0.01)	(p<0.01)	(p<0.05)	(p<0.05)	
diastolic pressure	composition of the invention	77±5	79±5	81±4	81±5	81±5
	equal volume of whole blood	9∓69	73±8	<b>∠∓9</b> ∠	<i>1</i> €∓7	82±12
•		(p<0.01)	(p<0.05)	(p<0.05)	(p<0.05)	
average arterial	composition of the invention	76±4	78±4	79±4	80±5	80±5
pressure	equal volume of whole blood	68±5	71±6	74±4	75±4	8∓08
		(p<0.01)	(p<0.01)	(p<0.05)	(p<0.05)	
cardiac	composition of the invention	77±5	79±5	81±4	81±4	81±5
contractivity	equal volume of whole blood	9∓69	73±8	76±7	249£	82±12
		(p<0.01)	(p<0.05)	(p<0.05)	(p<0.05)	
cardiac output		105±13	106±14	103±14	106±12	106±17
	equal volume of whole blood	$89\pm10$	83±18	83±15	88±23	$88\pm 16$
		(p<0.05)	(p<0.01)	(p<0.01)	(p<0.01)	
end-diastolic	composition of the invention	82±4	82±3	83±4	84±3	84±4
volume	equal volume of whole blood	78±7	78±9	80±7	82±8	84±8
		(p<0.05)	(p<0.05)			
				1000		

Comparison between composition of the invention and equal volume of whole blood For urine volume in restoration stage in shocked dogs [unit: ml/kg/h]

Table 2

		after transfusion	fusion	
	11	2h	3h	4h
Composition of the invention	2.23±1.03	0.94±0.22	0.95±0.29	1.00±0.30
equal volume of whole blood	$0.33\pm0.21$	$0.27\pm0.16$	$0.73\pm0.41$	$0.61\pm0.25$
	(p<0.01)	(p<0.01)		(p<0.05)

10

15

20

25

The composition of the Example 1 of the invention was administered to 48 patients in Hefei No. 105 Hospital, Anhui province. The total effective rate was 100%. Most patients, had the blood pressure raised, urine volume increased, and the limbs became warm during transfusion. Several patients whom the conventional drugs couldn't already reverse, the composition of the Example 1 of the invention begin to play its role 5-- 10minutes after infusion. The circulatory function of patients has recovered basically, (and there were no obvious clinical complications.

# Test 3. Experiment of acute toxicity

when dogs were given at 2.5 times the dosage for human, no adverse effects have been showed. At 5 times the recommended dosage, salivation and vomiting were seen in the dogs. At 3. 75 times the recommended dosage vomiting was seen without salivation in the dogs. All the above administered dogs survived more than 45 days. At 7. 5 times the recommended dosage the death occurred in the administered dogs. focal hemorrhage was seen in the lungs as target organs.

The composition of the present invention could be infused through vein at the dose of 8 ml of the composition of the present invention per kg body weight. It could be used directly in treating and saving the patients with shock, combined injuries or hematorrhea etc in order, to reverse the physiopathological status of patients and to obtain time for sequential treatment.

As compared with the prior art, the pharmaceutical composition of this invention has the following prominent features and improvement:

1. Greatly decreased volume of transfusion: In general, the dose for most patients is 500 ml or less than 500 ml. Even if the patients suffered from lethal hematorrhea, to infuse only 1/4 to 1/6 of volume of lost blood is enough. Thus, it could obviously decrease the incidence rate of tissue edema or overload of heart.

10

15

20

25

- 2. Rapid curative effect just during 5-- 10 minutes after infusion, the hemodynamics has been improved significantly.
- 3. Good efficacy. As Test 1 showed the composition of the invention had better efficacy than that of equal volume of fresh whole blood. Moreover, although the composition of the invention has no oxygen- carrying action, but it could improve micro--circulation and general status to decrease oxygen consumption and to increase oxygen transport. Thus at least 50% of blood transfused could be saved, it could mitigate the contradiction with short supply of blood, decrease the complications induced by blood transfusion, and reduce obviously the economic burden for the patients.
- 4. Maintain once of efficacy for long time. As Test 1 showed, after infusion of the composition of the invention, the improvement of hemodynamics and general condition could be maintained more than 3--4 hours, even if all other infusion and drugs were not administered.
- 5. Unnecessary special condition for storage: The composition could be stored at room temperature, simply used infused intravenously or intraosseously and conveniently transported, without special devices and special vehicles.
- 6. Unnecessary blood typing and cross match tests: It is suitable for individual with any blood type. Thus the valuable time could be gained to rescue the wounded and patients.
- 7. Wider uses: It could widely be used in the treatment of patients with shock of various types, brain trauma burn, combined injuries, cardiogenic shock induced by myocardial infarction of right ventricle, hypotension induced by hemodialysis, biliary pancreatitis, cardiovascular intoxication induced by narcotic, hepatic echinococcosis, and patients under operation.
- 8. Change of administration model: The composition of the invention could be infused drop by drop intravenously, instead of pushing so it could be

conveniently used with less complications.

In general, as compared with prior therapy, the composition of the present invention possesses unique benefit and theoretical innovation, for treating and saving the wounded and patients, so it is active significance.

10

### **Claims**

- 1. A pharmaceutical composition comprising 1.5--6.9 % (w/v) of one or more substances selected from sodium chloride, sodium bicarbonate, potassium chloride, magnesium sulfate, calcium chloride, calcium gluconate, calcium lactate, sodium lactate, sodium acetate and Tris (Hydroxy methyl) aminomethane, and 3~18 % (w/v) of one or more substances selected from hydroxyethylstarch, dextran, carboxymethylstarch, polyvinyl--pyrrolidone (PVP), gelatin derivatives, condensed glucose, glucose, fructose, lactose, glycerin, xylitol, sodium, alginate, N--2- hydroxypropylacrylamide, ethylene epoxide-polypropylene glycol, pectin, mannitol. and pentahydroxyethylstarch; as well as the remainder of conventional injections, as long as sodium chloride is not less than 1. 5 % (w/v), and the concentration of sodium ion is not more than that of 6.9 % (w/v) sodium chloride solution or equivalent.
- 2. The pharmaceutical composition of Claim 1, wherein the composition contains 4.2±0.2g sodium chloride and 7.6±0.6g hydroxyethylstarch per 100 ml.
- 3. The pharmaceutical composition of Claim 1 or 2, wherein the conventional injections are selected from water for injection, physiological saline, balanced buffers, glucose solution, sodium lactate solution, sodium acetate solution, Tris solution, and glucose and sodium chloride solution.
- 4. The pharmaceutical composition of Claim 1 or 2, wherein hydroxyethylstarch contains at least 10% hydroxyethylstarch with molecular weight of 25,000--45,000.
- 5. The pharmaceutical composition of Claim 1, wherein gelatin derivatives have molecular weight of 20,000~35,000, and are selected from urea--conjugated gelatin, modified liquid gelatin, oxidized polygelatin and

5

10

15

degraded gelatin polypeptide.

- 6. The pharmaceutical composition of Claim 1, wherein dextran has molecular weight of 40,000--230,000; carboxymethylstarch has molecular weight of 30,000--80,000, PVP has molecular weight of 5,000--700,000, condensed glucose has molecular weight of 8,000~ 12,000, sodium alginate has molecular weight of 20,000--26,000, pectin has molecular weight of 20,000-40,000; pentahydroxyethylstarch has molecular weight of 264,000.
- 7. A method for preparing the pharmaceutical composition of Claim 1, comprising: dissolving 3--18g of one or more substances selected from hydroxyethylstarch, dextran, carboxymethylstarch, PVP, gelatin derivatives, condensed glucose, glucose, fructose, lactose, glycerin, xylitol, sodium alginate, N--2--hydroxypropylacrylamide, ethylene epoxide-polypropylene glycol, pectin, mannitol, and pentahydroxyethylstarch, in total of 100 ml of one injection or mixture of several injections selected from water for injection, physiological saline, balanced buffers, glucose solution, sodium lactate solution, sodium acetate solution, Tris solution, and glucose and sodium chloride solution; the adding 1.5g sodium chloride and 0—5.4g of one or more substances selected from sodium chloride, sodium bicarbonate, potassium chloride, magnesium sulfate, calcium chloride, calcium gluconate, calcium lactate, sodium lactate, sodium acetate, and Tris with the proportion described above, then mixing, dissolving, to obtain the composition of the present invention.

10

### Abstract

The present invention relates to a pharmaceutical composition and the method for the preparation thereof, The composition comprises  $1.5\sim6.9\%$  (w/v) of one or more substances selected from sodium chloride, sodium bicarbonate, potassium chloride, magnesium sulfate, calcium chloride, calcium gluconate, and the like, and  $3\sim18\%$  (w/v) of one or more substances selected from hydroxyethylstarch, dextran, carboxymethylstarch, polyvinyl-pyrrolidone, gelatin derivatives, and the like as well as the remainder of conventional injections, as long as sodium chloride is not less than 1.5% (w/v). The pharmaceutical composition of the present invention is used to treat and save the wounded and patients, as well as to treat shock, its advantages include safe and convenient use, rapid and good curative effect, long time maintenance, extensive uses and the like.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

High Soft II Hom Who Live And Hall

i i į.

Approved for use through 6/30/98. OMB 0651-0032

Patent and Trademark Office; US DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number

									, , ,			
		DECLARA	TION - U	Itility	Or D	esign Pater	it Applicat	ion				
I hereby claim the benefit	under Title	35, United Stat	es Code §12	0 of ar	ıy Unıte	d States applic	ation(s) of an	y PCT	International appli	cation		
designating the United St	ates of Ameri	ica, listed belov	v and, insofa	r as the	s the subject matter of each of the claims of this application is not disclosed in the yield by the first paragraph of Title 35, United States Code §112, I acknowledge							
the duty to disclose inform	I internation:	al application if	the matter protection	provid	ed by th	e first paragrap	oh of Title 35,	United	States Code §112	, I acknowledge		
between the filing date of	the prior app	lication and the	e national or	as dell PCT i	nternatio	nue 37, Code (	of this applic	guiation	is § 1.36 which be	came available		
U.S Parent Appli			Parent	1011					Parant Datant	Numban		
Number	CHUON		ımber		Parent Filing Date (MM/DD/YYYY)				Parent Patent Number (if applicable)			
Transci		140	illibei		1 '	WIWI/DD/Y	111)		(іј арриса	101e)		
Additional U.S. or P												
As a named inventor	, I hereby a	appoint the f	ollowing r	egist	ered pi	ractitioner(s	) to prosecı	ute thi	s application a	nd to transact		
all business in the Pa	tent and Ti	rademark Of	fice conne	cted	therew	vith:						
	<b>区</b> Custo	omer Numb	er 20594									
		or					•		Place Custon	ner		
	☐ Regis	stered practit	tioner(s) n	ame/i	registr:	ation numbe	r listed hel	OW	Number Bar C	1		
		stered prueti	ioner(3) ii	ullic/	regiona	ation named	I listed ber	OW	Label Here	;		
		Dag	:-44:							.•		
NT			istration			3.7			Registra			
Name		N	umber	_	Name				Number			
		Į.		ł								
		İ										
		İ										
Additional registered p	ractitioner(s)	named on supp	lemental Re	gistere	d Practi	tioner Informa	tion sheet PTO	O/SB/0	2C attached hereto	).		
Direct all correspond	ence to 🗵	Customer	Number		2059	)4	or C	orres	ondence Addr	ess below		
•	or Bar Coo						-					
Name	Alvin R.				I			· · · · · ·				
Address	121111111											
City, State, Zip												
			1 1				1		1			
Country  Thereby dealers that all st		10	elephone				Fax					
I hereby declare that all sta be true; and further that the	uements mad ese statement	e nerein of my	own knowle	dge ar	e true ar	nd that all state	ments made o	on infor	mation and belief	are believed to		
imprisonment, or both, une	der Section 1	001 of Title 18	of the United	ieuge i i State	mat win es Code	and that such v	nents and the willful false st	nke so atemen	made are punisnat ts may jeonardize	the validity of		
the application or any pate	nt issued the	reon.				and man saon .	TITTAL TAIDO DE	atomici.	as may jeopardize	the varianty of		
Name of Sole or Firs	t Inventor	r	ΠAn	etitio	n has l	been filed fo	or this unsid	aned i	nventor			
		middle [if a		-	1							
	e (mst and	middle [ii a	пуј)				Family N	ame o	r Surname	<del> </del>		
Chaoying					ZHA	.0						
Inventor's Signature	9				L			Date	ad			
								Dau				
Residence: City	Shan	ghai	State			Country	P.R. Ch	ina	Citizenship	P.R. China		
Post Office Address	Chan	ahai Daad 1	70.7.602	<u> </u>						<u>.</u>		
1 ost Office Aduress	Chan	ghai Road 1	/0-/-602									
City	Shan	ghai	State	Т		Zip	200433		Country	P.R. China		
		_				-			·			
Additional inventor	s are being i	named on the	supp	oleme	ntal ado	ditional inven	itor(s) sheet(	(s) PTO	D/SB/02A attach	ed hereto.		
Name of Inventor			☐ A p	etitio	n has l	peen filed fo	r this unsig	gned in	nventor			
Given Name	e (first and	middle [if a	nvl)				Family No	ame o	r Surname			
	(**************************************	Alliadio [II a	)])				I diffilly 14	arric o	Guillaille			
Inventor's Signature	;							Date	ed			
Dosidones: Cit-			1 64 :	т		~						
Residence: City			State			Country			Citizenship			
Post Office Address			1		L				<u> </u>			
City			State			Zip			Country			